

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
 International Bureau



(43) International Publication Date
 2 October 2003 (02.10.2003)

PCT

(10) International Publication Number
WO 03/079816 A1

(51) International Patent Classification⁷: **A23L 1/212**,
 1/275, C09B 61/00, 67/54

(21) International Application Number: PCT/EP03/02749

(22) International Filing Date: 17 March 2003 (17.03.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
 MI2002A 000632 27 March 2002 (27.03.2002) IT

(71) Applicant (for all designated States except US): **INDENA S.P.A.** [IT/IT]; Viale Ortles, 12, 20139 Milano (IT).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **GIORI, Andrea** [IT/IT]; Viale Ortles 12, 20139 Milano (IT)

(74) Agents: **MINOJA, Fabrizio et al.**, Minoja Bracco Minoja S.r.l., Via Rossini 8, 20122 Milano (IT).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: A PROCESS FOR THE PREPARATION OF TOMATO EXTRACTS WITH HIGH CONTENT IN LYCOPENE

(57) Abstract: A process for the extraction of lycopene from whole tomatoes, in which process tomatoes are heat concentrated and extracted with water-saturated ethyl acetate.

WO 03/079816 A1

A PROCESS FOR THE PREPARATION OF TOMATO EXTRACTS WITH HIGH CONTENT IN LYCOPENE

FIELD OF THE INVENTION

The present invention relates to a process for the preparation of tomato extracts with high content in lycopene.

TECHNICAL BACKGROUND

5 Lycopene is a natural pigment, particularly abundant in tomatoes and watermelon, having intense red color. Due to this characteristic, as well as its safety and beneficial effects, lycopene is widely used in the food industry as a coloring agent, usually in the form of oleoresin, i.e. a suspension in natural lipids. In this form, lycopene oxidation (crystalline
10 lycopene being highly unstable) and bacterial degradation are prevented, ~~more~~ likely due to the lipids and natural antioxidants present. Furthermore, lycopene is used as food supplement thanks to its antioxidative and chemoprotective properties.

Although lycopene can be prepared by synthesis [Karrer *et al.*, *Helv.*
15 *Chim. Acta* 33, 1349 (1950); Isler *et al.*, *ibid.* 39, 463 (1956)], it is usually obtained by extraction from tomatoes (*Lycopersicum esculentum*). As lycopene has intense red color only when in the crystalline form, the extraction process should allow to obtain the product in this form.

To date, the suggested methods (WO 95/16363 and WO 97/48287)
20 comprise the separation of the serum from the pulp, and the extraction of the latter with solvents. In WO 97/48287 tomatoes, before pulp-serum separation, which has to be performed under controlled conditions, are subjected to heat treatment; the extraction is carried out in the hot as well.

Although these processes may be used with any type of tomatoes, the
25 lycopene starting content should preferably be above 50 ppm.

DETAILED DISCLOSURE OF THE INVENTION

The present invention relates to a process for the preparation of tomato whole extracts with lycopene content from 5% to 20% and with reducing sugars content, expressed as glucose, below 1%, which process comprises the following steps:

- a) pretreating fresh tomatoes, which comprises washing, then cutting or crushing;
- b) heat concentrating of the cut or crushed tomato from step a);
- c) extracting the concentrate from step b) with water-saturated ethyl acetate;
- d) backwashing the extract from step c) with water;
- e) concentrating the extract to dryness under reduced pressure.

Pre-treatment is carried out according to conventional techniques and any method providing a homogeneous cut/crushed tomato will be suitable.

Concentration (step b) is carried out by distillation under reduced pressure, at temperatures ranging from 40 to 70°C, preferably at 50°C, so that the weight of the cut/crushed tomato will be 20-30% the starting value.

Extraction of the concentrate (step c) is repeatedly performed with water-saturated ethyl acetate in a volume ranging from 1.0 to 2.5, preferably 2, times the weight of the concentrate, to obtain a lycopene-free residue. According to a preferred embodiment of the invention, the extraction is repeated four times. The extraction is carried out for at least one hour at room temperature, shielding from light and keeping the concentrate-solvent mixture under stirring.

Each extract is washed with water (step d), preferably in half the volume of the solvent used for each single extraction, after that the extracts are combined, filtered and evaporated to dryness under reduced pressure (step e). Washing with water is mandatory for the success of the process; it has in fact

been observed that, when this step is omitted, as illustrated in detail in the subsequent example 3, a higher amount of whole extract is obtained which has however percent lycopene content lower by about one third (approximately 4% instead of 6%, for tomatoes containing 50 ppm of lycopene).

5 Furthermore, the process of the invention allows to obtain crystalline lycopene, with purity higher than 50%, from which the oleoresin can be prepared. For this purpose, steps a)-d) are carried out as described above, whereas at step e) the extract is concentrated to a final volume ranging from 0.10 to 0.28% with respect to the starting volume. The concentrated extract is
10 then left to stand for some hours and the lycopene crystalline precipitate is filtered off and dried (step f). The resulting crystalline lycopene may optionally be suspended in ethanol, then filtered and washed with ethyl acetate until obtaining the desired purity. The oleoresin is obtained by adding seed oil to the lycopene crystals, preferably tomato seed oil or soybean oil (step g).

15 Advantageously, the process according to the present invention provides good yields even when using tomatoes with low starting content in lycopene and it allows to obtain a whole extract with high lycopene content, ranging from 5% to 20%, which is about twice higher than obtained with the method disclosed in WO 97/48287, as illustrated in the Comparison Example below.
20 This process is also advantageous in that the reducing sugars content in the extract is always lower than 1%, usually lower than 0.5%.

The invention is illustrated in greater detail by means of the following examples.

Example 1

25 **Preparation of the extract according to the invention**

52 kg of fresh tomatoes with lycopene content 50 ppm are cut and homogenized in a blender.

Part of water (34 L) is distilled off under reduced pressure (20 mBar) at

60°C and discarded, to obtain 17.8 Kg of tomato concentrate.

36 L of water-saturated ethyl acetate are poured on the concentrate and the mixture is stirred for 2 hours at room temperature, shielded from light.

After 2 hours the extract is collected and the residue is extracted again
5 with 36 L of water-saturated ethyl acetate. The mixture is stirred for 2 hours at room temperature, shielded from light. The extract is filtered and washed in a separatory funnel with 18 L of water, which is then removed and the extract is collected.

Two extractions and two washings as described above are repeated
10 (using 144 L of solvent totally). After filtration, the extracts are combined and concentrated to dryness under reduced pressure; the resulting tomato whole extract (38.9 g) has HPLC lycopene content of 6.05%, reducing sugars content (expressed as glucose) of 0.28%, phospholipids content of 12.97% and mono-
di- glycerids content of 24.02%.

15 Example 2

50 kg of fresh tomatoes with lycopene content of 150 ppm are cut and homogenized in a blender.

Part of water (31 L) is distilled off under reduced pressure (20 mBar) at 60°C and discarded, to obtain 18.8 Kg of tomato concentrate.

20 40 L of water-saturated ethyl acetate are poured on the concentrate and the mixture is stirred for 2 hours at room temperature, shielded from light.

After 2 hours the extract is collected and the residue is extracted again with 40 L of water-saturated ethyl acetate. The mixture is stirred for 2 hours at room temperature shielded from light. The extract is filtered and washed in a
25 separatory funnel with 20 L of water, which is then removed and the extract is collected.

Two extractions and two washings as described above are repeated (using 160 L of solvent totally). After filtration, the extracts are combined and

concentrated to dryness under reduced pressure; the resulting tomato whole extract (37.2 g) has HPLC lycopene content of 17.8% and reducing sugars content (expressed as glucose) of 0.31%.

Example 3

5 **Preparation of the extract without backwashing with water**

Tomatoes belonging to the same lot as in Example 1, with lycopene content of 50 ppm, are used.

4.5 kg of tomatoes are cut and homogenized in a blender, then 3.3 L of water are distilled off under reduced pressure (20 mBar) at 60°C.

10 The resulting concentrate (1.17 kg) is extracted 4 times with 2.3 L each of (9.2 L of solvent totally), stirring each time for 2 hours at room temperature and shielding from light.

The extracts are combined, filtered and concentrated to dryness under reduced pressure. The resulting whole extract (5.09 g) has HPLC lycopene
15 content of 4%, reducing sugars content (expressed as glucose) of 4.46%, phospholipids' content of 16.51% and a mono- di- glycerids content of 14.47%.

Example 4

Preparation of the oleoresin in tomato oil

The procedure of Example 1 is followed, but concentrating the combined
20 extracts to 200 ml final volume. The concentrated extract is left to stand overnight, shielded from light, to obtain a dark red needle crystal, which is filtered under vacuum, shielding from air, washed with ethyl acetate and dried under vacuum at 50°C, to obtain 4.23 g of crystalline lycopene with 51% purity.

25 The crystalline lycopene is added with 6.75 g of tomato seed oil (obtained by hexane extraction) and the mixture is stirred vigorously, to obtain 10.7 g of a fluid, homogeneous, dark red product having 19.8% lycopene content.

Example 5

Preparation of the oleoresin in soybean oil

The procedure of Example 1 is followed, but concentrating the combined extracts to a final volume of 200 ml. The concentrated extract is left to stand
5 overnight, shielded from light, to obtain a dark red needle crystal, which is filtered under vacuum, shielding from air, washed with ethyl acetate and dried under vacuum at 50°C, to obtain 4.23 g of crystalline lycopene with 51% purity.

The crystalline lycopene is added with 6.75 g of soybean oil (obtained by
10 hexane extraction) and the mixture is stirred vigorously, to obtain 10.7 g of a fluid, homogeneous, dark red product having 19.8% lycopene content.

Example 6

Preparation of 95% purity lycopene

The procedure of Example 1 is followed, but concentrating the combined
15 extracts to a final volume of 200 ml. The concentrated extract is left to stand overnight, shielded from light, to obtain a dark red needle crystal, which is filtered under vacuum, shielding from air. The solid is suspended in 80 ml of ethyl acetate and heated to 45°C with stirring for 20 min. The mixture is then left to cool to room temperature and filtered under vacuum, shielding from air.
20 The solid is suspended in 200 ml of ethanol and heated to 45°C with stirring for 10 min, then filtered while hot, under vacuum and shielding from air. This procedure is repeated once more. After that, the solid is washed on the filter with 40 ml of cold ethyl acetate, then dried under vacuum at 50°C, to obtain 2.05 g of crystalline lycopene with 95% purity.

Comparative Example

Extraction according to the method described in WO 95/16363

Tomatoes belonging to the same lot as in Example 1, with lycopene content of 50 ppm, are used.

5.14 kg of tomatoes are cut and homogenized in a blender, then centrifuged at 3000 r for 15', to separate serum from the insoluble fraction (1.315 kg), which is extracted 4 times with 2.65 L each of ethyl acetate (10.6 L of solvent totally), each extraction during 2 hours, under stirring, at a
5 temperature of 60°C and shielding from light.

The extracts are combined and concentrated to dryness under reduced pressure. The resulting tomato whole extract (6.07 g) has HPLC lycopene content of 3.5%, reducing sugars content (expressed as glucose) of 8.74%, phospholipids content of 35.57% and mono- di- glycerids content of 12.44%.

CLAIMS

- 1) A process for the preparation of tomato whole extracts with lycopene content from 5% to 20% and with reducing sugars content expressed as glucose lower than 1%, comprising the following steps:
- 5 a) pretreating fresh tomatoes, which comprises washing, then cutting or crushing;
- b) heat concentrating of the cut or crushed tomato from step a);
- c) extracting the concentrate from step b) with water-saturated ethyl acetate;
- 10 d) backwashing the extract from step c) with water;
- e) concentrating the extract to dryness under reduced pressure.
- 2) A process as claimed in claim 1, wherein the concentration of the extract according to step e) is carried out to a final volume ranging from 0.10 to 0.28% with respect to the starting volume, further comprising the following steps:
- 15 f) filtering and drying the lycopene precipitated from the concentrate; and optionally suspending lycopene in ethanol or ethyl acetate, then filtering and washing with ethyl acetate until obtaining the desired purity;
- 20 g) adding seed oil to lycopene from step f).
- 3) A process as claimed in claim 2, wherein the seed oil is tomato seed oil.
- 4) A process as claimed in claim 2, wherein the seed oil is soybean oil.
- 5) Tomato whole extracts with lycopene content from 5% to 20% and with content in reducing sugars, expressed as glucose, lower than 1%, obtainable
- 25 with the process of claim 1.
- 6) Crystalline lycopene with purity higher than 50% obtainable according to the process of claim 2-f).
- 7) Crystalline lycopene with purity higher than 90% obtainable according to

the process of claim 2-f).

8) Oleoresins containing lycopene of claim 7) obtainable with the process of any one of claims 2-4.

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/EP 03/02749

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A23L1/212 A23L1/275 C09B61/00 C09B67/54

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A23L C09B

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

WPI Data, EPO-Internal

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 97 48287 A (LYCORED NATURAL PROD IND LTD) 24 December 1997 (1997-12-24) cited in the application page 9, paragraph 2 -page 10, paragraph 1; claim 17; examples 1,2 ---	1,5,8
X	WO 96 13178 A (MAKHTESHIM CHEM WORKS LTD ;MAKHTESHIM AGAN OF NORTH AMERI (US); HA) 9 May 1996 (1996-05-09) the whole document ---	1-3,5-8
A	WO 95 16363 A (MAKHTESHIM CHEM WORKS LTD ;MAKHTESHIM AGAN OF NORTH AMERI (US); ZE) 22 June 1995 (1995-06-22) cited in the application claims; examples ----- -/--	1-8



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

Z document member of the same patent family

Date of the actual completion of the international search

15 July 2003

Date of mailing of the international search report

28/07/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Ketterer, M

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 03/02749

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 818 225 A (INDENA SPA) 14 January 1998 (1998-01-14) claims	1
X	example IV ---	6,7
A	DATABASE WPI Section Ch, Week 199916 Derwent Publications Ltd., London, GB; Class D21, AN 1999-186185 XP002247814 & JP 11 035444 A (NIKKO CHEM CO LTD), 9 February 1999 (1999-02-09) abstract	1
P,X	WO 02 072509 A (BASF AG ;JOHN MICHAEL (DE); WEGNER CHRISTOPH (DE)) 19 September 2002 (2002-09-19) examples ---	6,7
X	EP 1 103 579 A (NESTLE SA) 30 May 2001 (2001-05-30) column 2, line 46 - line 49; examples column 2, line 56 -column 3, line 7	6
A	column 1, line 46 -column 2, line 45 ---	1
E	WO 03 038064 A (SCHAAP ALBERT ;DSM NV (NL); AKISHINA RAISA ILLARIONOVNA (RU); VINE) 8 May 2003 (2003-05-08) page 26, line 21 - line 24 page 27, line 16 - line 17 ---	6,7
P,X	EP 1 201 762 A (VITATENE S A) 2 May 2002 (2002-05-02) page 5, line 20 - line 24 page 5, line 46 - line 48; claims 3,6; examples ---	6,7
X	DATABASE WPI Section Ch, Week 200032 Derwent Publications Ltd., London, GB; Class B04, AN 2000-373965 XP002247815 & RU 2 112 777 C (URALBIOFARM STOCK CO), 10 June 1998 (1998-06-10) abstract ---	8
X	EP 0 986 963 A (BASF AG) 22 March 2000 (2000-03-22) table 2 ---	6
X	US 5 858 700 A (AUSICH RODNEY L ET AL) 12 January 1999 (1999-01-12) examples ---	6-8

-/--

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 03/02749

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 3 467 579 A (BIANCHI MARIA LUISA ET AL) 16 September 1969 (1969-09-16) column 4, line 59 - line 64 ---	6,7
A	DATABASE WPI Section Ch, Week 200203 Derwent Publications Ltd., London, GB; Class E17, AN 2002-017919 XP002247816 & CN 1 298 904 A (SHENGMINGHONG SCI & TECHNOLOGY INVESTMEN), 13 June 2001 (2001-06-13) abstract -----	6-8

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 03/02749

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9748287	A	24-12-1997	IL 118697 A BR 9702303 A CN 1198661 A EP 0844831 A1 WO 9748287 A1 TR 9800268 T1	31-12-1999 02-03-1999 11-11-1998 03-06-1998 24-12-1997 22-06-1998
WO 9613178	A	09-05-1996	IL 111477 A AU 4104896 A CN 1176577 A , B DE 69530312 D1 EP 0789517 A1 JP 10509590 T WO 9613178 A1 US 5965183 A	14-07-1999 23-05-1996 18-03-1998 15-05-2003 20-08-1997 22-09-1998 09-05-1996 12-10-1999
WO 9516363	A	22-06-1995	IL 107999 A AU 690201 B2 AU 1513195 A WO 9516363 A1 US 5837311 A	08-02-1998 23-04-1998 03-07-1995 22-06-1995 17-11-1998
EP 0818225	A	14-01-1998	IT MI961442 A1 AT 227599 T AU 722774 B2 AU 2853997 A CA 2210039 A1 DE 69717015 D1 DK 818225 T3 EP 0818225 A1 ES 2186824 T3 JP 10226654 A PT 818225 T US 5897866 A	12-01-1998 15-11-2002 10-08-2000 29-01-1998 12-01-1998 19-12-2002 10-03-2003 14-01-1998 16-05-2003 25-08-1998 31-03-2003 27-04-1999
JP 11035444	A	09-02-1999	NONE	
WO 02072509	A	19-09-2002	DE 10103708 A1 WO 02072509 A1	01-08-2002 19-09-2002
EP 1103579	A	30-05-2001	EP 1103579 A1 AT 226613 T AU 7909200 A DE 69903645 D1 DE 69903645 T2 WO 0138443 A1 ES 2183471 T3 PT 1103579 T	30-05-2001 15-11-2002 04-06-2001 28-11-2002 13-03-2003 31-05-2001 16-03-2003 31-01-2003
WO 03038064	A	08-05-2003	WO 03038064 A2	08-05-2003
EP 1201762	A	02-05-2002	ES 2157166 A1 AU 6282600 A EP 1201762 A1 JP 2003507021 T CN 1370241 T WO 0112832 A1	01-08-2001 13-03-2001 02-05-2002 25-02-2003 18-09-2002 22-02-2001

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 03/02749

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
RU 2112777	C	10-06-1998	RU 2112777 C1	10-06-1998
EP 0986963	A	22-03-2000	DE 19841930 A1	16-03-2000
			CN 1252238 A	10-05-2000
			EP 0986963 A2	22-03-2000
			JP 2000103733 A	11-04-2000
			US 6235315 B1	22-05-2001
US 5858700	A	12-01-1999	AU 6056698 A	22-10-1998
			WO 9843620 A1	08-10-1998
US 3467579	A	16-09-1969	ES 343773 A1	01-12-1968
CN 1298904	A	13-06-2001	NONE	